

Reaction Mechanism Practice

STUDENT PRACTICE WORKSHEET • WORKSPACE PROVIDED

Name: _____ Date: _____

Directions: Select the best answer for each question. Show reasoning on the space provided.

1. 1-Bromobutane is treated with NaI in acetone. Which pathway is favored?

- A. SN1
- B. SN2
- C. E1
- D. E2

Reasoning / work:

2. 2-Bromobutane is heated with sodium ethoxide in ethanol. Which pathway is generally favored?

- A. SN1
- B. SN2
- C. E1
- D. E2

Reasoning / work:

3. 2-Bromobutane with NaN₃ in DMSO: which pathway is generally favored?

- A. SN2
- B. SN1
- C. E1
- D. Radical substitution

Reasoning / work:

4. tert-Butyl bromide with potassium tert-butoxide: which pathway is favored?

- A. SN2
- B. E2
- C. Electrophilic addition
- D. SN1 only

Reasoning / work:

5. Iodomethane and NaCN in DMSO: expected mechanism?

- A. SN1
- B. E1

C. SN2

D. E2

Reasoning / work:

6. Which elementary event occurs in a concerted E2 elimination?

- A. Leaving group departs before base acts
- B. Base removes beta H as leaving group leaves
- C. Carbocation rearranges first
- D. Nucleophile adds to alkene

Reasoning / work:

7. Which factor most strongly makes SN2 attack on neopentyl bromide unusually slow?

- A. Resonance of C–Br bond
- B. Beta branching creates steric hindrance
- C. Carbocation stability
- D. Aromaticity

Reasoning / work:

8. An SN2 reaction follows which idealized rate law?

- A. rate = $k[\text{RX}]$
- B. rate = $k[\text{Nu}]$
- C. rate = $k[\text{RX}][\text{Nu}]$
- D. rate = $k[\text{RX}]^2$

Reasoning / work:

9. For E2 on a cyclohexane chair, which arrangement is normally required?

- A. Leaving group equatorial and beta H equatorial
- B. Leaving group axial and beta H axial anti
- C. Leaving group and beta H on same carbon
- D. No beta hydrogen needed

Reasoning / work:

10. Which solvent often favors SN2 relative to the same reaction in a protic solvent?

- A. DMSO
- B. Water
- C. Methanol
- D. Ethanol

Reasoning / work:

11. At a stereogenic carbon, an SN2 reaction normally produces which geometric outcome?

- A. Retention
- B. Inversion
- C. Complete racemization
- D. No stereochemical change

Reasoning / work:

12. A planar carbocation is formed in an SN1 solvolysis. Which outcome is generally expected?

- A. Only inversion
- B. Only retention
- C. Loss of stereospecificity with both faces available
- D. No substitution

Reasoning / work:

13. For alkenes, E/Z labels depend on:

- A. Molecular mass only
- B. Highest CIP-priority substituent on each alkene carbon
- C. Hydrogen count alone
- D. Whether the alkene is terminal

Reasoning / work:

14. Two highest-priority groups on an alkene are on opposite sides. It is:

- A. E
- B. Z
- C. R
- D. S

Reasoning / work:

15. Two highest-priority groups on an alkene are on the same side. It is:

- A. E
- B. Z
- C. R
- D. S

Reasoning / work: