

Gas Laws — Practice Worksheet

Name: _____ Date: _____

Instructions: Show your reasoning, units, electron counts, or diagrams as needed. Use a separate sheet for additional work. For numeric results, use appropriate significant figures.

- 1. Ideal gas law: $n = 0.5$ mol; $P = 1.0$ atm; $T = 298.15$ K. Find volume (L).**

- 2. Ideal gas law: $n = 1.25$ mol; $V = 10.0$ L; $T = 300.15$ K. Find pressure (atm).**

- 3. Ideal gas law: $P = 0.95$ atm; $V = 5.0$ L; $T = 310.15$ K. Find moles (mol).**

- 4. Ideal gas law: $n = 2.0$ mol; $P = 2.5$ atm; $T = 273.15$ K. Find volume (L).**

- 5. Ideal gas law: $n = 0.75$ mol; $V = 12.0$ L; $T = 350.0$ K. Find pressure (atm).**

- 6. Ideal gas law: $P = 1.25$ atm; $V = 2.5$ L; $T = 298.15$ K. Find moles (mol).**

7. Boyle law: $P_1 = 1.2 \text{ atm}$; $V_1 = 3.5 \text{ L}$; $P_2 = 0.8 \text{ atm}$. Find $V_2 \text{ (L)}$.

8. Boyle law: $P_1 = 0.95 \text{ atm}$; $V_1 = 8.0 \text{ L}$; $P_2 = 1.9 \text{ atm}$. Find $V_2 \text{ (L)}$.

9. Charles law: $V_1 = 2.4 \text{ L}$; $T_1 = 300.0 \text{ K}$; $T_2 = 360.0 \text{ K}$. Find $V_2 \text{ (L)}$.

10. Charles law: $V_1 = 5.0 \text{ L}$; $T_1 = 250.0 \text{ K}$; $T_2 = 300.0 \text{ K}$. Find $V_2 \text{ (L)}$.

11. Gay Lussac law: $P_1 = 1.1 \text{ atm}$; $T_1 = 280.0 \text{ K}$; $T_2 = 350.0 \text{ K}$. Find $P_2 \text{ (atm)}$.

12. Gay Lussac law: $P_1 = 2.0 \text{ atm}$; $T_1 = 320.0 \text{ K}$; $T_2 = 256.0 \text{ K}$. Find $P_2 \text{ (atm)}$.