

Spectroscopy & Structure Practice

STUDENT PRACTICE WORKSHEET • WORKSPACE PROVIDED

Name: _____ Date: _____

Directions: Select the best answer for each question. Show reasoning on the space provided.

1. A strong IR absorption near 1715 cm^{-1} is consistent with:

- A. C=O stretch in a simple saturated ketone
- B. $\text{C}\equiv\text{N}$ stretch
- C. O–H stretch only
- D. C–H bending only

Reasoning / work:

2. A broad IR band spanning roughly $3200\text{--}3600\text{ cm}^{-1}$ often indicates:

- A. Alcohol O–H
- B. Nitrile $\text{C}\equiv\text{N}$
- C. Alkyne $\text{C}\equiv\text{C}$ only
- D. Carbonyl C=O

Reasoning / work:

3. A carboxylic acid O–H stretch is often characterized by:

- A. Sharp peak at 2250 cm^{-1}
- B. Very broad absorption about $2500\text{--}3300\text{ cm}^{-1}$
- C. Single sharp peak at 1600 cm^{-1}
- D. No IR absorption

Reasoning / work:

4. A relatively sharp IR absorption around 2250 cm^{-1} can indicate:

- A. $\text{C}\equiv\text{N}$
- B. C–O single bond
- C. Alcohol O–H
- D. Alkane C–H

Reasoning / work:

5. An aldehydic ^1H NMR signal commonly appears near:

- A. 0–1 ppm
- B. 2–3 ppm
- C. 9–10 ppm

D. 14–16 ppm

Reasoning / work:

6. An aromatic proton often resonates in approximately which ^1H NMR region?

- A. 0–1 ppm
- B. 6.5–8.5 ppm
- C. 12–16 ppm
- D. 18–20 ppm

Reasoning / work:

7. An ethyl fragment often shows what integration and splitting pattern in first-order ^1H NMR?

- A. 3H triplet and 2H quartet
- B. 3H singlet and 3H singlet
- C. 1H septet and 6H doublet
- D. 2H singlet and 2H singlet

Reasoning / work:

8. A 6H doublet plus 1H septet can support which fragment?

- A. Methoxy group
- B. Isopropyl group
- C. Ethene
- D. Aldehyde

Reasoning / work:

9. A TLC spot travels 2.4 cm while the solvent front travels 6.0 cm. R_f = ?

- A. 0.25
- B. 0.40
- C. 2.5
- D. 3.6

Reasoning / work:

10. In simple acetone, the two methyl groups give what expected ^1H NMR pattern?

- A. Two 3H doublets
- B. One 6H singlet
- C. One 3H triplet
- D. One 1H aldehyde singlet

Reasoning / work:

11. A curved arrow in an organic mechanism starts at:

- A. The electron source
- B. The atom receiving electrons only
- C. The positive charge always
- D. The reaction arrow

Reasoning / work:

12. Lower pKa generally indicates:

- A. Stronger acid
- B. Weaker acid
- C. Stronger base only
- D. No connection to acidity

Reasoning / work:

13. Which is a Lewis base?

- A. Electron-pair donor
- B. Electron-pair acceptor
- C. Only a proton donor
- D. A compound with no electrons

Reasoning / work:

14. What is the degree of unsaturation for C₆H₁₂O (neutral closed-shell)?

- A. 0
- B. 1
- C. 2
- D. 3

Reasoning / work:

15. What is the degree of unsaturation for C₆H₆?

- A. 2
- B. 3
- C. 4
- D. 5

Reasoning / work: