

Epoxide Ring Opening: Acidic vs Basic Conditions

STUDENT PRACTICE WORKSHEET

Name: _____ Date: _____

Instructions: Select the best answer for questions 1–8 and briefly justify your choice using substrate, reagents, or reaction conditions. For 9–10, explain your reasoning in complete sentences.

1. Basic nucleophile attack on propylene oxide usually occurs at:

- A. Less substituted carbon B. More substituted carbon C. Epoxide oxygen D. No carbon

Reasoning: _____

2. When one epoxide carbon is tertiary, acid-catalyzed opening often favors attack at:

- A. More substituted carbon B. Less substituted carbon exclusively C. No carbon D. O atom only

Reasoning: _____

3. After hydrolysis/workup, the epoxide oxygen typically becomes:

- A. An alcohol OH B. A ketone O automatically C. CO
- ₂
- D. An ether absent any nucleophile

Reasoning: _____

4. A basic epoxide opening is generally:

- A. SN
- ₂
- like and backside B. SN
- ₁
- with free long-lived carbocation C. E
- ₂
- always D. Radical chain

Reasoning: _____

5. Propylene oxide + NaOCH₃/MeOH predominantly gives:

- A. CH
- ₃
- CH(OH)CH
- ₂
- OCH
- ₃
- B. CH
- ₃
- CH(OCH
- ₃
-)CH
- ₂
- OH C. CH
- ₃
- COCH
- ₃
- D. CH
- ₃
- CH
- ₂
- CH
- ₃

Reasoning: _____

6. 2-Methyl-1,2-epoxypropane + MeOH/H⁺ often gives:

- A. (CH
- ₃
-)
- ₂
- C(OCH
- ₃
-)CH
- ₂
- OH B. (CH
- ₃
-)
- ₂
- C(OH)CH
- ₂
- OCH
- ₃
- C. 2-Methylpropene D. Acetone

Reasoning: _____

7. Epoxide ring opening is driven in part by:

- A. Relief of ring strain B. Creation of a more strained ring C. Loss of all oxygen D. An aromaticity gain

Reasoning: _____

8. Stereochemical attack at the reacting epoxide carbon tends to:

- A. Invert configuration B. Retain via front-side attack always C. Erase stereochemistry at all centers D. Give only radicals

Reasoning: _____

9. Compare basic and acid-catalyzed methanol opening of 2-methyl-1,2-epoxypropane.

10. Does protonated epoxide opening always involve a freely rearranging carbocation?
