

INTOCHEMISTRY

Stoichiometry Practice Problems

ANSWER KEY

Solutions include units and setups.

Use: H 1.01; C 12.01; N 14.01; O 16.00; Ca 40.08; Fe 55.85 g/mol. Avogadro's number = 6.022×10^{23} mol⁻¹.

1. Balance the equation by writing the smallest whole-number coefficients: $__ \text{H}_2 + __ \text{O}_2 \rightarrow __ \text{H}_2\text{O}$

Solution: $2 \text{H}_2 + 1 \text{O}_2 \rightarrow 2 \text{H}_2\text{O}$. Coefficients: 2, 1, 2. Hydrogen and oxygen atoms are conserved.

2. How many moles of H_2 are required to react completely with 4.00 mol O_2 ? Use the equation in Question 1.

Solution: $4.00 \text{ mol O}_2 \times (2 \text{ mol H}_2 / 1 \text{ mol O}_2) = 8.00 \text{ mol H}_2$.

3. $\text{CH}_4 + 2 \text{O}_2 \rightarrow \text{CO}_2 + 2 \text{H}_2\text{O}$. How many moles of CO_2 form from 2.50 mol CH_4 (excess O_2)?

Solution: $2.50 \text{ mol CH}_4 \times (1 \text{ mol CO}_2 / 1 \text{ mol CH}_4) = 2.50 \text{ mol CO}_2$.

4. $4 \text{Fe} + 3 \text{O}_2 \rightarrow 2 \text{Fe}_2\text{O}_3$. How many moles of Fe_2O_3 form from 3.00 mol O_2 (excess Fe)?

Solution: $3.00 \text{ mol O}_2 \times (2 \text{ mol Fe}_2\text{O}_3 / 3 \text{ mol O}_2) = 2.00 \text{ mol Fe}_2\text{O}_3$.

5. $\text{N}_2 + 3 \text{H}_2 \rightarrow 2 \text{NH}_3$. How many moles of NH_3 form from 7.50 mol H_2 (excess N_2)?

Solution: $7.50 \text{ mol H}_2 \times (2 \text{ mol NH}_3 / 3 \text{ mol H}_2) = 5.00 \text{ mol NH}_3$.

6. $2 \text{H}_2 + \text{O}_2 \rightarrow 2 \text{H}_2\text{O}$. What mass of water forms from 5.00 g H_2 (excess O_2)?

Solution: $M(\text{H}_2) = 2.02 \text{ g/mol}$; $M(\text{H}_2\text{O}) = 18.02 \text{ g/mol}$. $5.00 \text{ g H}_2 \times (1 \text{ mol} / 2.02 \text{ g}) \times (2 \text{ mol H}_2\text{O} / 2 \text{ mol H}_2) \times (18.02 \text{ g/mol}) = 44.6 \text{ g H}_2\text{O}$.

7. $\text{CH}_4 + 2 \text{O}_2 \rightarrow \text{CO}_2 + 2 \text{H}_2\text{O}$. What mass of CO_2 forms from 8.00 g CH_4 (excess O_2)?

Solution: $M(\text{CH}_4) = 16.05 \text{ g/mol}$; $M(\text{CO}_2) = 44.01 \text{ g/mol}$. $8.00 \text{ g CH}_4 \times (1 \text{ mol} / 16.05 \text{ g}) \times (1 \text{ mol CO}_2 / 1 \text{ mol CH}_4) \times (44.01 \text{ g/mol}) = 21.9 \text{ g CO}_2$.

8. $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$. What mass of CaO forms from 12.5 g CaCO_3 ?

Solution: $M(\text{CaCO}_3) = 100.09 \text{ g/mol}$; $M(\text{CaO}) = 56.08 \text{ g/mol}$. $12.5 \text{ g} \times (1 \text{ mol} / 100.09 \text{ g}) \times (1 \text{ mol CaO} / 1 \text{ mol CaCO}_3) \times (56.08 \text{ g/mol}) = 7.00 \text{ g CaO}$.

9. $2 \text{H}_2 + \text{O}_2 \rightarrow 2 \text{H}_2\text{O}$. A mixture starts with 3.00 mol H_2 and 2.00 mol O_2 . Identify the limiting reactant, find moles of H_2O made, and moles of excess reactant remaining.

Solution: 3.00 mol H_2 needs 1.50 mol O_2 . H_2 is limiting; 3.00 mol H_2O form; $2.00 - 1.50 = 0.500$ mol O_2 remains.

10. $\text{N}_2 + 3 \text{H}_2 \rightarrow 2 \text{NH}_3$. A mixture starts with 2.00 mol N_2 and 5.00 mol H_2 . Identify the limiting reactant and calculate moles of NH_3 formed.

Solution: 2.00 mol N_2 requires 6.00 mol H_2 , but only 5.00 is available. H_2 limits. $5.00 \times (2/3) = 3.33 \text{ mol NH}_3$.

11. The theoretical yield of CO_2 is 8.00 g. An experiment recovers 6.40 g. Calculate percent yield.

Solution: $(6.40 \text{ g} / 8.00 \text{ g}) \times 100\% = 80.0\%$.

12. How many moles of water molecules are present in 1.80×10^{24} molecules of H_2O ?

Solution: $(1.80 \times 10^{24} \text{ molecules}) \div (6.022 \times 10^{23} \text{ molecules/mol}) = 2.99 \text{ mol H}_2\text{O}$.